



TITLE:

Measurement of methane flux over an evergreen coniferous forest canopy using a relaxed eddy accumulation system with tuneable diode laser spectroscopy detection

AUTHOR(S):

Sakabe, Ayaka; Hamotani, Ken; Kosugi, Yoshiko; Ueyama, Masahito; Takahashi, Kenshi; Kanazawa, Akito; Itoh, Masayuki

CITATION:

Sakabe, Ayaka ...[et al]. Measurement of methane flux over an evergreen coniferous forest canopy using a relaxed eddy accumulation system with tuneable diode laser spectroscopy detection. Theoretical and Applied Climatology 2012, 109(1-2): 39-49

ISSUE DATE:

2012-07

URL:

<http://hdl.handle.net/2433/157358>

RIGHT:

The final publication is available at www.springerlink.com; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。

- 1 Measurement of methane flux over an evergreen coniferous forest canopy using a relaxed eddy
2 accumulation system with tuneable diode laser spectroscopy detection
3
4 Corresponding author: Ayaka Sakabe,
5 e-mail address: sakabea@kais.kyoto-u.ac.jp
6 Tel.: 81-75-753-6149
7 Fax: and fax numbers: 81-75-753-6149
8
9
10 Ayaka Sakabe,¹ Ken Hamotani,² Yoshiko Kosugi,¹ Masahito Ueyama,² Kenshi Takahashi,³ Akito
11 Kanazawa,¹ Masayuki Itoh⁴
12 ¹Laboratory of Forest Hydrology, Graduate School of Agriculture, Kyoto University, Kitashirakawa
13 Oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan
14 ² School of Life and Environmental Sciences, Osaka Prefecture University, Gakuen-cho 1-1, Sakai,
15 599-8531, Japan
16 ³ Research Institute for Sustainable Humanosphere, Kyoto University, Gokasho, Uji , 611-0011, Japan
17 ⁴Center for Ecological Research, Kyoto University, Hirano, Otsu, 520-2113, Japan
18

19

20 **Abstract**

21 Very few studies have conducted long-term observations of methane (CH_4) flux over forest canopies. In

22 this study, we continuously measured CH_4 fluxes over an evergreen coniferous (Japanese cypress) forest

23 canopy throughout 1 year, using a micrometeorological relaxed eddy accumulation (REA) system with

24 tuneable diode laser spectroscopy (TDLS) detection. The Japanese cypress forest, which is a common

25 forest type in warm-temperate Asian monsoon regions with a wet summer, switched seasonally between a

26 sink and source of CH_4 probably because of competition by methanogens and methanotrophs, which are

27 both influenced by soil conditions (e.g., soil temperature and soil moisture). At hourly to daily timescales,

28 the CH_4 fluxes were sensitive to rainfall, probably because CH_4 emission increased and/or absorption

29 decreased during and after rainfall. The observed canopy-scale fluxes showed complex behaviours

30 beyond those expected from previous plot-scale measurements and the CH_4 fluxes changed from sink to

31 source and *vice versa*.

33 **1. Introduction**

34 Most research on methane (CH_4) fluxes from natural ecosystems has focused on wetlands or rice

35 paddy fields, which are thought to be large CH_4 sources and thus important components of the global CH_4

36 budget. Considering that a major portion of forest soil is water-unsaturated, forests are generally assumed

to be an insignificant atmospheric CH₄ sink, representing to about 6% of the global sink (Le Mer and Roger 2001). However, studies have revealed that forest ecosystems, especially in wet warm climates, are not always CH₄ sinks. For example, some upland soils could be a CH₄ source when soils become water-saturated following precipitation events (Silver et al. 1999; Savage et al. 1997; van den Pol-van Dasselaar et al. 1998). Wang and Bettany (1997) also reported that upland soils that were incubated anaerobically began producing CH₄ within days or weeks. However, the CH₄ dynamics in whole forest ecosystems are still poorly understood because of insufficient information on topographically complex landscapes. It is important to quantify CH₄ fluxes in forest ecosystems because they cover a large portion of continental areas.

Long-term CH₄ flux measurements in forested areas have been mostly performed using chamber methods. While chamber methods are useful for understanding the processes controlling CH₄ fluxes on small spatial scales (usually less than 1 m²), the small footprint of the measurement creates a difficult scaling problem (Denmead 1994) when estimating landscape-scale fluxes in heterogeneous terrain such as forests. Moreover, the occasional measurements of manual chambers restrict the time resolution. These inherent limitations of chamber methods have made it difficult to evaluate CH₄ dynamics in whole forest ecosystems, as CH₄ fluxes from forest soils have wide spatial and temporal variations. CH₄ fluxes in forest ecosystems could have wide-ranging spatio-temporal variations both emission and absorption sides, especially in the forest ecosystems which have wide spatio-temporal range in soil water status, such as

55 Asian monsoon forests under warm and humid climate, or boreal and tropical peat forests. Scaling up of
56 CH₄ fluxes in those types of forests and understanding CH₄ dynamics as a whole ecosystem would be
57 difficult only with chamber methods. In addition, both open and closed chambers disturb natural
58 environmental conditions during the measurement by affecting airflow, radiant energy receipt, and energy
59 transfer to the atmosphere (Denmead 1994). Consequently, CH₄ exchanges estimated by the chamber
60 method could contain biases and uncertainties if up-scaled to stand, watershed, and regional scales.

61 Micrometeorological methods such as the eddy covariance (EC) method are ideally suited for
62 continuous canopy-scale flux measurements integrated over a larger area without artificial disturbance.
63 Although the EC method has been widely used for canopy-scale flux measurements, it has not been used
64 until very recently for CH₄ measurements because it requires a fast-response and high-precision gas
65 analyzer. Recent technological advances in the application of tuneable diode laser spectroscopy (TDLS)
66 to *in situ* field measurements open the possibility of long-term EC measurement of CH₄ in a variety of
67 ecosystems. Previously, a limited number of CH₄ EC measurements have been obtained in peatlands
68 (Hendriks et al. 2008; Schrier-Uijl et al. 2009), rice paddy fields (Simpson et al. 1994), and prairies (Kim
69 et al. 1998a, b). The lack of long-term CH₄ flux observations in forest ecosystems restricts our
70 understanding of canopy-scale CH₄ dynamics. However, measuring CH₄ exchange over forest ecosystems
71 is still challenging compared to measurements in the above source areas because of the small fluxes in
72 forests (Smeets et al. 2009). According to previous EC measurements in wetlands and farmlands, the

precision of the CH₄ concentration measurements was 2.9 ppb at 10 Hz using a Quantum Cascade Laser Spectroscopy (QCLS) analyzer (QCL-TILDAS-76; Aerodyne Research Inc., Billerica MA, USA) (Kroon et al. 2007). Recently, the open path sensor is available for the CH₄ EC measurement with the precision of < 5 ppb at 10 Hz (McDermitt et al. 2011). However, those precisions are still insufficient to measure small CH₄ fluxes at forest ecosystems if the CH₄ fluxes measured by chamber techniques (Itoh et al. 2005, 2007, 2009) assumed to be representative to the canopy scale exchange.

Although the state-of-the-art CH₄ analyzers could be insufficient for the EC measurements, those analyzers are available for CH₄ flux measurements by using a micrometeorological relaxed eddy accumulation (REA) method (Businger and Oncley 1990; Hamotani et al. 1996, 2001). The REA method can take longer time for CH₄ concentration measurements, thus a laser signal can be averaged to optimize the instrumental sensitivity, and higher precision of TDLS CH₄ analyzer was achievable. The flux, calculated by the REA method, is equal to the difference in the mean concentrations of the trace gas of interest associated with updraft and downdraft, multiplied by the standard deviation of the vertical wind velocity and an empirical coefficient.

In this study, we employed an REA method (Businger and Oncley 1990; Hamotani et al. 1996, 2001) with a TDLS CH₄ analyzer for long-term observation of CH₄ fluxes from a temperate evergreen coniferous forest site. Our goal was to examine whether the REA method is applicable to (1) measure CH₄ fluxes over the forest canopy, (2) reveal the amplitude and seasonal variations in CH₄ fluxes, and (3)

examine the response of CH₄ fluxes to rainfall. This is the first report showing the seasonal cycle of canopy-scale CH₄ fluxes in a temperate forest.

2. Methods

2.1 Site description

The observations were made in a coniferous forest in the Kiryu Experimental Watershed, (KEW; area: 5.99 ha) in Shiga Prefecture, Japan. A meteorological observation tower is located in a small catchment within KEW (Fig. 1). The forest consists of 50-year-old Japanese cypress (*Chamaecyparis obtusa* Sieb. et Zucc.). Mean tree height was approximately 16.8 m in 2007. The annual mean air temperature and precipitation measured at the meteorological station shown in Fig. 1 from 2002 to 2009 were 13.3°C and 1,576 mm yr⁻¹, respectively. The site has a warm temperate monsoon climate with a wet summer. Rainfall occurs throughout the year with two peaks in summer due to the Asian monsoon; the early summer ‘Baiu’ front season and the late summer typhoon seasons. The entire watershed is underlain by weathered granite with considerable amounts of albite.

Canopy fluxes of heat, water, and CO₂ have also been measured at this site at 29-m above the ground, using the EC method (e.g., Takanashi et al. 2005; Kosugi et al. 2007; Kosugi and Katsuyama 2007; Ohkubo et al. 2007). Takanashi et al. (2005) reported that the CO₂ flux for 92% of the daytime flux and 81% of the nighttime flux originated from the forest area according to an analytical footprint model by

Schuepp et al. (1990). The trend in wind direction at this site did not change seasonally but had diurnal variations. The daytime wind direction was from all directions, whereas the night-time wind direction was mainly from the south (Kosugi et al. 2007). Some wetlands were located in riparian zones along streams within the flux footprint area, which were either always submerged or periodically submerged. The streams and the main wetland areas (approximately 10^0 - 10^2 m²) are shown in Fig. 1. The riparian zones and wetlands were distributed in both the north and south directions within the flux footprint area. The size of these areas could slightly increase after rainfall. Notably, an express highway was opened approximately 400 m south of the tower in February of 2008. During night-time, this highway was always inside of the flux footprint. The paddy fields were situated several km north to west of the tower, although these area were mostly out of the flux footprint (Takanashi et al. 2005). CH₄ fluxes from wetlands and water-unsaturated soils were investigated using a chamber method with a gas chromatograph analyzer (Itoh et al. 2005, 2007, 2009). Comparisons of the CH₄ fluxes from the previous chamber data and the present REA data will be described later.

2.2. Measurements

The CH₄ flux was measured using the REA method (Businger and Oncley 1990; Hamotani et al. 1996, 2001). Although the REA method is theoretically the same as the EC method, it does not require a fast response from the gas analyzer compared to the EC method. Compared to wetlands, forest canopies are

more challenging sites for conducting CH₄ flux measurements by the EC method. This is because the emission and absorption of CH₄ are both relatively small over forest canopies, and TDLS analyzers may not always be precise enough to detect the fluxes. The greatest benefit of applying the REA method is that sufficient precision in CH₄ concentration TDLS measurements is achievable by signal averaging over a longer duration. The precision of the TDLS CH₄ analyzer (FMA-200; Los Gatos Research, Mountain View, CA, USA) is 3 ppb at 10 Hz rate, 1 ppb at 1 Hz rate and 0.1 ppb at 100 s rate according to the catalog specifications. Moreover, the REA method can save electronic consumption and minimize the possible noises induced by the pressure drift of the measurement cell of the TDLS CH₄ analyzer, because the REA method with lower sampling frequency can measure CH₄ fluxes without a high-power vacuum pump. The another advantage of the REA method is that it does not require any corrections such as high frequency attenuation for closed-path EC method or WPL correction (Webb et al. 1980) for open-path EC methods, which could obscure the observed small flux values at the forest ecosystems. One possible disadvantage of the REA method involves the switching speed of the valve system, which may lead to the loss of high frequency information, however those effects can be negligible at measurements over tall forest canopies, such as in our forest (Ueyama et al. 2009). Equation 1 expresses CH₄ flux (F_{CH_4} , nmol m⁻² s⁻¹) (Subsequently, we use this abbreviation only for CH₄ flux measured by the REA method). This equation can also be used to calculate CO₂ flux (F_{CO_2} , μmol m⁻² s⁻¹):

$$F_{CH_4} = \sigma_w \left(\overline{S_{CH_4}^+} - \overline{S_{CH_4}^-} \right) \rho_a b \quad (1)$$

146

147 where σ_w is the standard deviation (SD) of the vertical wind velocity (w , m s⁻¹), $\overline{S_{CH_4}^+}$ and $\overline{S_{CH_4}^-}$
148 are the 30-minute mean CH₄ mole fractions (ppmv) associated with updraft and downdraft, respectively,
149 and ρ_a (mol m⁻³) is molar air density. The coefficient b was empirically determined from EC data using air
150 temperature given by Eq. 2:

151

$$b = \frac{\overline{w'T'}}{\sigma_w (\overline{T^+} - \overline{T^-})} \quad (2)$$

153

154 where $\overline{T^+}$ and $\overline{T^-}$ are 30-min mean fast-response temperature data associated with updraft and
155 downdraft, respectively. We determined b to be 0.59, which is the average of 16,576 values obtained from
156 Eq. 2 and the standard error was 0.04. The sensible heat flux and air temperature data used here was all
157 data measured from August 2009 to August 2010 by the sonic anemometer (SAT-550; Kaijo Corp., Tokyo,
158 Japan) mounted on top of the 29-m-tall tower. The term b was applied to both F_{CO_2} and F_{CH_4} . The value of
159 b is relatively constant over a wide range of atmospheric stability (Bowling et al. 1998).

160 Our REA system consists of a sonic anemometer (SAT-550; Kaijo Corp.) to measure wind speed and
161 direction, two diaphragm pumps (CV-201, Enomoto Co., Tokyo, Japan), four reservoirs (CCK-20; GL
162 Science, Tokyo, Japan) to accumulate sampled gas, a CO₂/H₂O gas analyzer (LI-840; Li-Cor Inc., Lincoln,

NE, USA), a CH₄ gas analyzer (Baer et al. 2002; Hendriks et al. 2008) (FMA-200; Los Gatos Research) and a data logger (CR1000; Campbell Scientific, Logan, UT, USA). The sonic anemometer was mounted on top of the 29-m-tall tower and the air inlets were set directly below the sonic anemometer. Air samples for updraft and downdraft were pulled through DK tubes (inner tube: 4 mm in diameter and coated with aluminium) to the reservoirs by two diaphragm pumps at a constant flow rate (0.7 l min⁻¹). The frequency of switching the pumps was 10 Hz. One pump worked only for updraft and the other worked only for downdraft. Updraft or downdraft was determined by the difference between the instantaneous and adjacent 15-min moving average of *w*. Air flow was switched using the solenoid valves (CKD USB3-6-3-E; CKD Corp., Aichi, Japan) and controlled by the CR1000 data logger. To sequentially determine the flux every half hour, we prepared two sets of sampling reservoirs: one pair of reservoirs accumulated air during 0–30 min and the other pair accumulated air for the next 30–60 min. After accumulating in the reservoirs for 30 min, the air in the reservoirs was pulled a diaphragm pump into the CH₄ analyzer at a flow rate of approximately 0.7 l min⁻¹, and the air in each reservoir was analyzed for 2 min. Three filters were inserted in the gas sample line to protect the CH₄ analyzer from dust and insects. Before entering the CO₂/H₂O and the CH₄ analyzers, the sampled air was dried using a gas dryer (PD-50T-48; Perma Pure Inc., Toms River, NJ, USA). Dilution by water vapour, which could not be completely removed by the drying system, was corrected for using the H₂O concentration measured with the CO₂/H₂O analyzer. We confirmed that the gas dryer did not alter the CH₄ mixing ratio within

measurement uncertainties. Data were recorded at 10 Hz by the CR1000 data logger and stored on a compact flash card using a compact flash module (CFM100, Campbell Scientific). A 10-s moving average filtered the high frequency noises for the CO₂/H₂O analyzer and a 1-s moving average was used for the CH₄ analyzer. Volumetric soil water content (VSWC) at a depth of 0–30 cm was measured with a CS616 water content reflectometer (Campbell Scientific) at four different points around the tower on the water-unsaturated forest floor, and soil temperatures were measured with a thermistor (RT-10,11,12; ESPEC Mic Corp., Kanagawa, Japan) at depths of 2 cm near the tower. Precipitation was measured with a tipping bucket rain gauge at an open screen site near the tower. Air temperature above the canopy was measured with a ventilated temperature and humidity sensor (HHP45AC; Vaisala, Helsinki, Finland) at a height of 29 m above the ground.

In this study, we compared CO₂ fluxes measured by the REA and EC methods in order to examine the validity of our REA system. An open-path CO₂/H₂O analyzer (LI-7500; Li-Cor) was used to measure CO₂ flux by the EC method. The double-rotation method was applied to the sonic anemometer velocities (McMillen 1988), and the Webb, Pearman, Leuning (WPL) correction for the effect of air density fluctuations (Webb et al. 1980) was applied to CO₂ flux using the EC method. Details of the EC measurements have been described by Kosugi et al. (2007) and Okubo et al. (2007b).

2.3. Data analysis

199 Hendriks et al. (2008) described the specifications of the CH₄ analyzer used in this study in detail. To
200 examine the accuracy and precision of our CH₄ analyzer, calibration experiments were performed on site
201 using a standard CH₄ gas cylinder (Takachiho, Tokyo, Japan, 1773 ppb CH₄ in synthetic air) several times
202 during the course of this study. The typical SD for determining the CH₄ mixing ratio was 0.4 ppb with a
203 30-s moving average within a 3-min standard gas flow period, which is the same condition used for
204 calculating the reservoirs concentrations with the REA method; the reservoir concentrations were
205 averaged for 30 s and analyzed within 2 min. No significant drift in the measurement accuracy of the CH₄
206 analyzer was observed during the entire observation period (less than 6 ppb). We also examined the
207 detection limit of CO₂ and CH₄ fluxes in our REA system by storing the same air in reservoir pairs and
208 measuring the concentration difference in each pair. This check mode was performed during 1 day of
209 every month.

210 We performed a *t*-test (significance level: 0.05) for the CH₄ concentration difference between updraft
211 and downdraft to examine whether there was a statistical difference between the mean values. F_{CH_4}
212 assumed to be zero by the *t*-test (21.8% of all available data) is shown as grey circles in Figs. 3 and 5. All
213 CO₂ and CH₄ fluxes collected with the REA method were rejected when CO₂ flux data collected with the
214 EC method did not meet the stationary criteria (Foken and Wichur 1996; Aubinet et al. 2000). We also
215 rejected CO₂ and CH₄ fluxes collected during night-time under highly stratified conditions, using a
216 previously examined friction velocity threshold of 0.3 m s⁻¹ for the CO₂ flux data (Takanashi et al. 2005).

217 The total amount of F_{CH_4} data rejected by these criteria accounted for 66% of the entire data series.

218 The data analyzed in this study were recorded between 1 August 2009 and 31 August 2010. Data were
219 missing from 11 to 20 August 2009, from 16 June to 22 July 2010, and from 10 December 2009 to 17
220 February 2010 due to instrumental malfunctions.

221

222 3. Results

223 3.1. Validity of the CH_4 fluxes collected with the REA system

224 Before application to F_{CH_4} measurement, we validated our REA system by comparing F_{CO_2} measured
225 by the EC and REA method for daytime (Fig. 2a) and nighttime periods (Fig. 2b). The CO_2 fluxes by the
226 EC and REA methods were highly correlated for both daytime with a slope of 0.95, $r^2 = 0.72$ and the root
227 mean square error (RMSE) of $3.7 \mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 2a) and nighttime with a slope of 0.81, $r^2 = 0.45$ and
228 the RMSE of $3.4 \mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 2b). Although the slopes of the linear regression showed a slightly
229 smaller value than 1.0, the observed data both in the daytime and nighttime were mostly scattered around
230 the 1:1 line and the F_{CO_2} measured by the REA method did not show obviously higher or lower values
231 than those by the EC method (Fig. 2a, b). The F_{CO_2} measured by the EC and the REA methods had worse
232 correlation in the night-time than daytime (Fig. 2b).

233 The F_{CH_4} detection limit obtained from the check mode with our REA system showed a diurnal
234 variation because it depended on turbulent intensity (i.e., σ_w in Eq. 1 is larger in the daytime than in the

night-time) The night-time (0:00–6:00 and 18:00–24:00) and daytime (6:00–18:00) F_{CH_4} detection limits averaged for all 14 check mode days were 4.2 ± 3.7 and $7.4 \pm 5.9 \text{ nmol m}^{-2} \text{ s}^{-1}$, respectively, and the detection limits did not change seasonally.

3.2. Amplitude and seasonal variations in CH_4 flux and its response to rainfall

Figure. 3 shows the seasonal variations in (a) instantaneous F_{CH_4} , (b) air and soil temperature, and (c) precipitation and VSWC. The average and SD of F_{CH_4} was $5.9 \pm 11.5 \text{ nmol m}^{-2} \text{ s}^{-1}$ for the summer of 2009 (August and September), $5.3 \pm 10.4 \text{ nmol m}^{-2} \text{ s}^{-1}$ for the fall of 2009 (October and November), $2.2 \pm 10.9 \text{ nmol m}^{-2} \text{ s}^{-1}$ for the winter of 2009 (December, February, and March), $-10.0 \pm 14.3 \text{ nmol m}^{-2} \text{ s}^{-1}$ for the spring of 2010 (April, May, and June), and $-4.7 \pm 15.3 \text{ nmol m}^{-2} \text{ s}^{-1}$ for the summer of 2010 (July and August). This site had a heterogeneous topography, and some riparian zones and wetlands were distributed both in the north and south directions within the flux footprint area. However, the wind direction at this site did not change seasonally. We confirmed that both daytime and night-time F_{CH_4} from the north or south was not particularly larger than those from other wind directions by analysing F_{CH_4} for each wind direction (Fig. 4). F_{CH_4} seasonally shifted from emission in the summer and fall of 2009 to absorption in the spring of 2010. Then the absorption weakened and changed to emission in the summer of 2010 (Fig. 3a). The diurnal patterns changed seasonally. In the summer and fall of 2009, F_{CH_4} showed clear diurnal variation with an emission peak around noon (Figs. 5a, b and 6a, b). Large emission was

observed in the fall of 2009 during sequential rain events for several days (Fig. 5b). The emission decreased with a decrease in air temperature (Fig. 6a, b), and then F_{CH_4} became almost zero in winter (Figs. 5c and 6c). F_{CH_4} remained relatively small until the spring of 2010 (Fig. 5d). Then, F_{CH_4} gradually shifted to exhibit a clear diurnal variation with an absorption peak around noon, which was an opposite pattern compared to the previous summer (Fig. 5e and 6d). The magnitude of peak CH_4 absorption increased with air temperature. Maximum CH_4 absorption was observed in June, when the absence of rain lasted 17 days and VSWC decreased. After intense rainfalls in late July 2010, the CH_4 absorption rate gradually decreased and seemed to shift to emission.

We detected evidence for short-term F_{CH_4} that rainfall was an important factor contributing to increased CH_4 emission. After rainfall on 12 September 2009, high F_{CH_4} was observed on 13 September 2009 (Fig. 5a) and similarly on 27 October and 2 November 2009 (Fig. 5b). Averaged diurnal variations for the summer of 2009 (August and September) and the fall of 2009 (October and November) also showed that CH_4 emission increased after precipitation (Fig. 6a, b). Even when high CH_4 absorption rates were observed around noon, rainfall contributed to weakening CH_4 absorption and/or F_{CH_4} switched to emission as shown on 24 May 2010 (Fig. 5e). Averaged diurnal variation for the spring of 2010 (April, May, and June) also showed that CH_4 absorption was weakened after precipitation (Fig. 6d). Approximately a day after rainfall, CH_4 emission typically increased and/or absorption decreased (Fig. 5a, b, e). The response of F_{CH_4} to rainfall was not obvious in winter (Figs. 5c and 6c).

4. Discussion

4.1. Assignment of the REA system

From comparison of CO₂ fluxes between the REA and EC data, we concluded that our REA system generally provided a good approximation of the EC method and could thus be applied to measure F_{CH_4} . One possible reason for the difference in CO₂ fluxes between the EC and REA methods is the method for determining coefficient b in Eq. (1); coefficient b , although constant in Eq. 1, changes slightly and is not constant throughout the day or year. Another possibility is that the loss of high frequency information due to the switching speed of the REA valve system might lead to an underestimation of F_{CO_2} with the REA method. However, the latter possibility was thought to be less likely because turbulent transfer was significant at a lower frequency below 1 Hz at this site (Kosugi et al. 2007; Ueyama et al. 2009).

The most difficult part of measuring F_{CH_4} in the forest canopy is that the F_{CH_4} detection limit is very close to the actual flux range. Although F_{CH_4} was larger than the detection limit in summer, fall, and spring, it was smaller in winter. Chamber studies revealed that F_{CH_4} ranged from approximately 0 to 720 nmol m⁻² s⁻¹ in wetlands and from -1.7 to 1.4 nmol m⁻² s⁻¹ in water-unsaturated forest floors (Itoh et al. 2005, 2007, 2009); some of these values are smaller than the detection limit of our REA system. We suggest that further improvements in measurement precision for the difference between upward and downward air CH₄ concentration are required to measure F_{CH_4} more precisely particularly in the winter. A

possible approach to improving the precision of our REA system is to apply the hyperbolic REA method (Bowling et al. 1999), which samples air only for high turbulent transport eddies. Applying this method would allow us to increase the CH_4 concentration differences (up to a factor of 2.7; Bowling et al. 1999) and thus improve the precision of the REA system. One disadvantage of the hyperbolic REA is that the majority of air (80%) is discarded, and roughly 10% of the original volume is sampled into each of the updraft and downdraft reservoirs (Bowling et al. 2003). The information contained in the discarded air must be reconstructed through the coefficient b . The b is determined under the assumption of the scalar similarity between temperature, CO_2 and CH_4 , so under some conditions large errors can result from violation of scalar similarity using the hyperbolic REA (Ruppert 2002).

4.2. Amplitude and seasonal variations in CH_4 flux and its response to rainfall

At the study site, the main CH_4 sources areas (riparian wetlands) were heterogeneously distributed within the flux footprint area, as shown in Fig. 1, and the typical wind direction (described in Section 2.1) did not change seasonally. Although there were large CH_4 sources in the north and south, daytime and night-time F_{CH_4} from the north or south were not particularly larger than those from other wind directions (Fig. 4). Although there was a difference between the F_{CH_4} from the north and south in summer 2010, this was probably caused by artificial difference due to limited number of available data; F_{CH_4} in summer 2010 could not be affected by wind direction. Thus, we assumed that the observed F_{CH_4} seasonal

variations would be mainly caused by the activity of methanogens and methanotrophs influenced by the soil temperature and water conditions but not wind direction.

Itoh et al. (2005, 2007, 2009) measured seasonal CH₄ flux variations from different parts of the slope in a water-unsaturated forest floor and riparian wetlands at the study site using the chamber method from 2001 to 2005. They showed that CH₄ emission rates from wetlands ranged from 0 to 720 nmol m⁻² s⁻¹, and that the emission rates increased significantly during high soil temperature and VSWC periods. They hypothesised that the emission rates were large enough to turn the entire watershed into a net CH₄ source, even though the source areas were very limited (Itoh et al. 2005). The emission rates from the wetlands varied from year to year due to the hydrological conditions that changed in relation to precipitation patterns (particularly summer precipitation patterns) (Itoh et al. 2007). The reported CH₄ fluxes from the water-unsaturated forest floor at this site ranged from -1.7 to 1.4 nmol m⁻² s⁻¹ (Itoh et al. 2009). CH₄ uptake was usually observed throughout the sampling periods; however, on lower hillslopes, where groundwater constantly existed underground, weak CH₄ uptake was observed only at low soil temperature (< 15°C); these areas turned to a CH₄ source at high soil temperatures (Itoh et al. 2009). The chamber measurements revealed that CH₄ fluxes in this forest were heterogeneous at both temporal and spatial scales, thus highlighting the importance of conducting continuous measurements of canopy-scale CH₄ fluxes integrated over a larger area to evaluate the total CH₄ budget.

Four new insights were obtained by combining the results from earlier chamber studies and this study.

325 First, the canopy-scale F_{CH_4} results by the REA method showed emission in the summer and fall of 2009,
326 indicating that a forest ecosystem consisting mostly of a water-unsaturated forest floor could be a CH_4
327 source for an entire watershed, possibly due to a large source contribution by a tiny wetland (Whalen et al.
328 1990; Keller and Reiners 1994; Hudgens and Yavitt et al. 1997; Itoh et al. 2007). Another possible source
329 was the contribution from the lower hillslopes of the water-unsaturated forest floor. Even though the soil
330 surface was water-unsaturated, groundwater constantly existed underground and produced CH_4 under
331 anaerobic conditions, which could have been emitted (Itoh et al. 2009). This study shows, for an entire
332 watershed, that a Japanese cypress forest in a warm temperate climate could be a CH_4 source in summer
333 and fall according to tower based measurements. Previous chamber-based studies speculated on the
334 canopy-scale CH_4 dynamics from the plot-scale measurements, which might contain large uncertainties.
335 Our canopy-scale measurements support the results of earlier chamber measurements. We found that
336 monthly averaged F_{CH_4} in October of 2009 was $6.6 \text{ nmol m}^{-2} \text{ s}^{-1}$ and it was not far less than the average
337 CH_4 fluxes in a peat meadow ($29.7 \text{ nmol m}^{-2} \text{ s}^{-1}$, Hendriks et al. 2008), a dairy farm ($42.7 \text{ nmol m}^{-2} \text{ s}^{-1}$,
338 Kroon et al. 2007) and a boreal fen ($80 \text{ nmol m}^{-2} \text{ s}^{-1}$, Long et al. 2010). It is not negligible CH_4 emission
339 from forest ecosystems, and it is important to quantify CH_4 fluxes in forest ecosystems and monitor it in
340 the long-term.

341 Second, continuous measurements at high temporal resolution revealed that the canopy-scale CH_4
342 emissions increased and/or absorption decreased after rainfall particularly in summer and fall of 2009.

343 The F_{CH_4} response to rainfall was caused by changes in the soil water condition because the methanogenic
344 activities increase and methanotrophic activities decrease in anoxic environments (Le Mer and Roger
345 2001). The area of the CH_4 source was broadened along the riparian zones after rainfall, and the lower
346 hillslope part of the water-unsaturated forest floor may have switched from a sink to a source for CH_4
347 because the anaerobic area may also have broadened deep underground, as mentioned above. Another
348 possibility is that CH_4 diffusion from the air to the soil may have been inhibited by rainfall causing a
349 decline in CH_4 absorption (Bradford et al. 2001). As shown in Fig. 5a, b, e, the influence of rainfall to
350 F_{CH_4} was obvious 1 or 2 days after rainfall events, at the short time-scale. On the other hand, F_{CH_4} tended
351 to be influenced by rainfall in the seasonal time-scale as seen in the shift of F_{CH_4} to emission during
352 summer 2010 (Fig. 3a). F_{CH_4} seemed to shift from absorption to emission during about 2 weeks after
353 intense rainfalls in late July 2010 (Fig. 3a). These different time-scale responses of F_{CH_4} to rainfall were
354 probably caused by different processes, and should be investigated with more data. In winter, the
355 response of F_{CH_4} to rainfall appeared to be low, suggesting that less CH_4 was produced and/or absorbed in
356 winter compared to other seasons; the activities of both methanogens and methanotrophs are low at low
357 soil temperatures, although methanotrophs are much less temperature dependent (Dunfield et al. 1993).

358 Third, CH_4 absorption rates increased from spring 2010 as soil temperatures increased and VSWC
359 decreased. These results are consistent with previous chamber measurements (Itoh et al. 2009). Uptake
360 rates increased as soil temperature increased in the lower range of temperature ($<15^\circ\text{C}$) and as VSWC

decreased. Previous studies reported a similar activation of methanotrophs with an increase in temperature (Whalen et al. 1990; Dobbie et al. 1996; Prime and Christensen 1997; Ishizuka et al. 2000). In contrast, methanogens function at intermediate temperature ranges from 20 to 40°C (Yamane et al. 1961), and their activity is extremely low at low temperatures (Dunfield et al. 1993). The large F_{CH_4} absorption obtained in the spring of 2010 by the REA method was possibly due to the different responses of methanogens and methanotrophs to temperature. Methanotrophs might function well under conditions where methanogens are still unable to function well (i.e., low temperature) (Dunfield et al. 1993; La Mer and Roger 2001). We still do not have enough information to explain the differences in F_{CH_4} between fall 2009 and spring 2010 within a similar temperature range. These differences represent complex behaviour beyond that expected on the basis of previous plot-scale measurements. The difference in temperature before the sampling season (summer or winter in this case) might have affected both methanogenic and methanotrophic activities during these periods. Longer duration observational data are needed to help clarify the seasonal F_{CH_4} variation in forest ecosystems.

Finally, the CH_4 absorption rates in spring 2010 (monthly averaged F_{CH_4} from April to June of 2010 ranged from -5.1 to -18.3 $nmol\ m^{-2}\ s^{-1}$) were larger than those reported by the previous chamber study (-1.7 to 1.4 $nmol\ m^{-2}\ s^{-1}$; Itoh et al. 2009) at the water-unsaturated forest floor at this site. Although the CH_4 fluxes obtained by the REA and chamber measurements cannot be compared directly, one possible explanation involves the detection limit of our REA system ($4.2 \pm 3.7\ nmol\ m^{-2}\ s^{-1}$ in the night-time and

379 $7.4 \pm 5.9 \text{ nmol m}^{-2} \text{ s}^{-1}$ in the daytime), which may not be sufficient to detect the range of CH_4 uptake rates
380 obtained by the chamber method for the water-unsaturated forest floor (-1.7 to $1.4 \text{ nmol m}^{-2} \text{ s}^{-1}$). The
381 other possible reason is that the chamber method is not always an excellent tool for investigating
382 representative CH_4 flux over a large watershed. An unobserved area that is consuming CH_4 more
383 effectively than estimated previously might exist at the study site. Recently, we have been conducting
384 continuous chamber measurements of CH_4 flux using the same TDLS CH_4 analyzer at the
385 water-unsaturated forest floor at the study site and have found several times higher CH_4 absorption rates
386 than the earlier chamber measurements at one of three automated chambers (unpublished data). These
387 results suggest that canopy-scale measurements using micrometeorological methods are important to
388 evaluate total CH_4 exchange and its impact on the environment.

389

390 **Conclusions**

391 This study is the first report of continuous measurements of canopy-scale F_{CH_4} in an upland forest
392 using the REA method with a TDLS CH_4 analyzer. Our observations revealed how the entire forest
393 ecosystem complexly switched between being a CH_4 sink or a source on hourly, diurnal, and seasonal
394 scales. As micrometeorological methods provide spatially integrated fluxes with high temporal
395 resolutions, measuring F_{CH_4} with these methods is particularly important to investigate the CH_4 dynamics
396 in forest ecosystems. In this study, we demonstrated that the REA method is applicable for measuring

CH₄ flux over a large representative area. Further longer-term observations with improvement of the system are needed to evaluate the controlling factors for methanogens and methanotrophs activity in the CH₄ dynamics of forest ecosystems. A combined approach between conventional chamber and micrometeorological methods is particularly important to evaluate the total CH₄ exchange, its impact on the environment, and the detailed processes involved.

Acknowledgements

This work was supported in part by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan.

Figure captions

Figure 1 Topographic map of the observation site. Precipitation was recorded at the indicated meteorological station. The streams are shown as bold lines, and riparian zones are located along the streams. The main wetland areas are in black.

Figure 2 Comparison between CO₂ fluxes obtained by the REA and EC methods at the half-hourly time scale in the **a** daytime (0600-1800 h) and **b** nighttime (0000-0600 and 1800-2400 h). The thin and bold lines represent 1:1 and linear regression, respectively. The linear regression equation and r^2 obtained from analysis of all data are shown.

415 Figure 3 Seasonal variation in **a** instantaneous canopy CH₄ flux and CH₄ flux assumed to be zero by the
416 *t*-test (grey circles), **b** air temperature and soil temperature (grey lines), and **c** precipitation and
417 the volumetric soil water content (VSWC) at the water-unsaturated forest floor during August
418 2009 and August 2010, in an evergreen Japanese cypress forest in a warm temperature climate.

419 Figure 4 Daytime and night-time CH₄ fluxes averaged for each wind direction (north, south, east, west) in
420 each season: **a** the sum of all seasons, **b** summer 2009, **c** fall 2009, **d** winter 2009, **e** spring 2010,
421 and **f** summer 2010 are shown. Daytime and night-time CH₄ fluxes are shown as white and
422 black circles. Error bars show the standard deviations. The data used for averaging seasonal CH₄
423 fluxes for each wind direction are shown as bars. The data for daytime and night-time are shown
424 as white and black bars, respectively.

425 Figure 5 Diurnal course of instantaneous canopy CH₄ flux and CH₄ flux assumed to be zero by the *t*-test
426 (grey circles), precipitation, and air temperature and soil temperature (dotted lines) measured in
427 **a** summer (between 5 September and 14 September 2009), **b** fall (between 25 October and 3
428 November 2009), **c** winter (between 4 December and 8 December 2009 and between 7 March
429 and 11 March 2010), **d** early spring (between 1 March and 5 March 2010), and **e** late spring
430 (between 21 May and 30 May 2010) in an evergreen Japanese cypress forest in a warm
431 temperature climate.

432 Figure 6 Diurnal variations of CH₄ fluxes before precipitation (black circle) and after precipitation (white

433 circle) (in case accumulated precipitation was more than 2 mm half hour⁻¹) and air temperature,
434 averaged for **a** the summer of 2009 (August and September), **b** the fall of 2009 (October and
435 November), **c** the winter of 2009 (December, February and March), and **d** the spring of 2010
436 (April, May and June). Before and after precipitation were discriminated according to
437 accumulated precipitation within 24 h was more than 2 mm half hour⁻¹ or not. Error bars show
438 the standard deviations for CH₄ fluxes.

439

440 **References**

- 441 Aubinet M, Grelle A., Ibrom A, Rannik U, Moncrieff J, Foken T, Kowalski AS, Martin PH, Berbigier P,
442 Bernhofer C, Clement R, Elbers J, Granier A, Grunwald T, Morgenstern K, Pilegaard K, Rebmann
443 C, Snijders W, Valentini R, Vesala T (2000) Estimates of the annual net carbon and water exchange
444 of forests: the EUROFLUX methodology. Adv. Ecol. Res. 30:113–175.
- 445 Baer DS, Paul JB, Gupta M, O’Keefe A. (2002) Sensitive absorption measurements in the near-infrared
446 region using off-axis integrated cavity-output spectroscopy. Appl. Phys. B75:261–265.
- 447 Bowling DR, Pataki DE., Ehleringer JR (2003) Critical evaluation of micrometeorological methods for
448 measuring ecosystem-atmosphere isotopic exchange of CO₂. Agric. Meteorol. 3118:1-21.
- 449 Bowling DR, Delany AC, Turnipseed AA, Baldocchi DD, Monson RK (1999) Modification of the
450 relaxed eddy accumulation technique to maximize measured scalar mixing ratio differences in

- 451 updrafts and downdrafts. *J. Geophys. Res.* 104:9121-9133.
- 452 Bowling DR, Turnipseed AA, Delany AC, Baldocchi DD, Greenberg JP, Monson RK (1998) The use of
- 453 relaxed eddy accumulation to measure biosphere-atmosphere exchange of isoprene and other
- 454 biological trace gases. *Oecologia.* 116:306-315.
- 455 Bradford MA, Ineson P, Wookey PA, Lappin-Scott HM (2001) Role of CH₄ oxidation, production and
- 456 transport in forest soil CH₄ flux. *Soil Biol. and Biochem.* 33:1625-1631.
- 457 Businger JA, Oncley SP (1990) Flux measurement with conditional sampling. *J. Atmos. Oceanic Technol.*
- 458 7:349-352.
- 459 Denmead OT (1994) Measuring fluxes of CH₄ and N₂O between agricultural systems and the atmosphere.
- 460 In: Peng S, Ingram KT, Neue H-U, Ziska LH (eds) *Climate change and rice*. International Rice
- 461 Research Institute, Manila, Philippines.
- 462 Dobbie KE, Smith KA, Christensen S, Degorska A, Orlanski P (1996) Effect of land use on the rate of
- 463 methane uptake by surface soils in Northern Europe. *Atmos., Environ.* 30:1005-1011.
- 464 Dunfield P, Knowles R, Dumont R, Moore TR (1993) Methane production and consumption in
- 465 temperate and subarctic peat soils: response to temperature and pH. *Soil Biol. Biochem.*
- 466 25:321-326.
- 467 Foken T, Wichura B (1996) Tools for quality assessment of surface-based flux measurements. *Agric. For.*
- 468 *Meteorol.* 78:83-105.

- 469 Hamotani K, Uchida Y, Monji N, Miyata A (1996) A system of the relaxed eddy accumulation method to
470 evaluate CO₂ flux over plant canopies. *J. Agric. Meteorol.* 52(2):135–139.
- 471 Hamotani K, Monji N, Yamaguchi K (2001) Development of a long-term CO₂ flux measurement system
472 using REA method with density correction. *J. Agric. Meteorol.* 57(2):93–99.
- 473 Hendriks DMD, Dolman AJ, van der Molen MK, van Huissteden J (2008) A compact and stable eddy
474 covariance set-up for methane measurements using off-axis integrated cavity output spectroscopy.
475 *Atmos. Chem. Phys.* 8:431–443.
- 476 Hudgens DE, Yavitt JB (1997) Land-use effects on soil methane and carbon dioxide fluxes in near Ithaca,
477 New York. *Ecoscience.* 4(2):214–222.
- 478 Ishizuka S, Sakata T, Ishizuka K (2000) Methane oxidation in Japanese forest soils. *Soil Biol. Biochem.*
479 32:769–777.
- 480 Itoh M, Ohte N, Katsuyama M, Koba K, Kawasaki M, Tani M (2005) Temporal and spatial variability of
481 methane flux in a temperate forest watershed. *J. Japan Soc. Hydrol. Water Resour.* 18:244–256. (in
482 Japanese with English abstract)
- 483 Itoh M, Ohte N, Koba K, Katsuyama M, Hayamizu K, Tani M (2007) Hydrologic effects on methane
484 dynamics in riparian wetlands in a temperate Forest catchment. *J. Geophys. Res.* 112(G1):G01019.
- 485 Itoh M, Ohte N, Koba K (2009) Methane flux characteristics in forest soils under an East Asian monsoon
486 climate. *Soil Biol. Biochem.* 41:388–395.


- 487 Keller M, Reiners WA (1994) Soil-atmosphere exchange of nitrous oxide, nitric oxide, and methane
488 under secondary succession of pasture to forest in the Atlantic lowlands of Costa Rica. *Global*
489 *Biogeochem. Cycles*. 8(4):99-409.
- 490 Kim J, Verma SB, Billesbach DP (1998a) Seasonal variation in methane emission from a temperate
491 *Phragmites*-dominated marsh: effect of growth stage and plant-mediated transport. *Global Change*
492 *Biol*. 5:433–440.
- 493 Kim J, Verma SB, Billesbach DP, Clement RJ (1998b) Diel variation in methane emission from a
494 midlatitude prairie wetland: significance of convective throughflow in *Phragmites australis*. *J.*
495 *Geophys. Res.* 103:28,029–28,039.
- 496 Kosugi Y, Katsuyama M (2007) Evapotranspiration over a Japanese cypress forest. II. Comparison of the
497 eddy covariance and water budget methods. *J. Hydrol.*, 334:305–311.
- 498 Kosugi Y, Takanashi S, Tanaka H, Ohkubo S, Tani M, Yano M, Katayama T (2007) Evapotranspiration
499 over a Japanese cypress forest. I. Eddy covariance fluxes and surface conductance characteristics
500 for 3 years. *J. Hydrol.* 337:269–283.
- 501 Kroon PS, Hensen A, Jonker HJJ, Zahniser MS, van't Veen WH, Vermeulen AT (2007) Suitability of
502 quantum cascade laser spectroscopy for CH₄ and N₂O eddy covariance flux measurements.
503 *Biogeosciences*. 4:715-728.
- 504 LeMer J, Roger P (2001) Production, oxidation, emission and consumption of methane by soils: a review.

- 505 Eur. J. of Soil Biol. 37:25–50.
- 506 Long KD, Franagan LB, Cai T (2010) Diurnal and seasonal variation in methane emissions in a northern
- 507 Canadian peatland measured by eddy covariance. Global Change Biol. 16:2420–2435.
- 508 McDermitt D, Burba G, Xu L, Anderson T, Komissarov A, Riensche B, Schedlbauer J, Starr G, Zona D,
- 509 Oechel W, Oberbauer S, Hastings S (2011) A new low-power, open-path instrument for measuring
- 510 methane flux by eddy covariance. Appl Phys. B102:391-405.
- 511 McMillen RT, (1988) An eddy correlation technique with extended applicability to non-simple terrain.
- 512 Boundary-Layer Meteorol. 43:231–245.
- 513 Ohkubo S, Kosugi Y (2007) Amplitude and seasonality of storage fluxes for CO₂, heat and water vapor in
- 514 a temperate Japanese cypress forest. Tellus. 60B:11–20.
- 515 Ohkubo S, Kosugi Y, Takanashi S, Mitani T, Tani M (2007) Comparison of the eddy covariance and
- 516 automated closed chamber methods for evaluating nocturnal CO₂ exchange in a Japanese cypress
- 517 forest. Agric. For. Meteorol. 142:50–65.
- 518 Prime A, Christensen S (1997) Seasonal and spatial variation of methane oxidation in a Danish spruce
- 519 forest. Soil Biol. Biochem. 29:1165-1172.
- 520 Ruppert J (2002) Eddy sampling methods for the measurement of trace gas fluxes, Diploma thesis, 95pp,
- 521 University of Bayreuth.
- 522 Savage K, Moore TR, Crill PM (1997) Methane and carbon dioxide exchanges between the atmosphere

- 523 and northern boreal forest soils. *J. Geophys. Res.* 102:29279–29288.
- 524 Schrier-Uijl AP, Veenendaal EM, Leffelaar PA, van Huissteden JC, Berendse F (2009) Methane
- 525 emissions in two drained peat agro-ecosystems with high and low agricultural intensity. *Plant Soil.*
- 526 doi:10.1007/s11104-009-0180-1.
- 527 Schuepp PH, Leclerc MY, MacPherson JJ, Desjardins RL (1990) Footprint prediction of scalar fluxes
- 528 from analytical solutions of the diffusion equation. *Boundary-Layer Meteorol.* 50:335–373.
- 529 Silver WL, Lugo AE, Keller M (1999) Soil oxygen availability and biogeochemistry along rainfall and
- 530 topographic gradients in upland wet tropical forests soil. *Biogeochemistry.* 44:301–328.
- 531 Simpson IJ, Thurtell GW, Kidd GE, Lin M, Demetriades-Shah TH, Flitcroft ID, Kanemasu ET, Nie D,
- 532 Bronson KF, Neue HU (1994) Tunable diode laser measurements of methane fluxes from an
- 533 irrigated rice paddy field in the Philippines. *J. Geophys. Res.* 100:7283–7290.
- 534 Smeets CJPP, Holzinger R, Vigano I, Goldstein AH, Röckmann T (2009) Eddy covariance methane
- 535 measurements at a Ponderosa pine plantation in California. *Atmos. Chem. Phys.* 9:8365–8375.
- 536 Takanashi S, Kosugi Y, Tanaka Y, Yano M, Katayama T, Tanaka H, Tani M (2005) CO₂ exchange in a
- 537 temperate Japanese cypress forest compared to that in a cool-temperate deciduous broadleaved
- 538 forest. *Ecol. Res.* 20:313–324.
- 539 Ueyama M, Hamotani K, Nishimura W (2009) A technique for high-accuracy flux measurement using a
- 540 relaxed eddy accumulation system with an appropriate averaging strategy. *J. Agric. Meteorol.*

- 541 65(4):315–325.
- 542 Van den Pol-van Dasselaar A, Van Beusichem ML, Oenema O (1998) Effects of soil moisture content
- 543 and temperature on methane uptake by grasslands on sandy soils. *Plant Soil*. 204:213-222.
- 544 Wang FL, Bettany JR (1997) Methane emission from Canadian prairie and forest soils under short term
- 545 flooding conditions. *Nutr. Cycling Agroecosyst*. 49:197–202.
- 546 Webb EK, Pearman GI, Leuning R (1980) Correction of flux measurements for density effects due to heat
- 547 and water vapour transfer. *Q., J. R. Meteorol.* 106:85–100.
- 548 Whalen SC, Reeburgh WS, Sandbeck KA (1990) Rapid methane oxidation in a landfill cover soil. *Appl*
- 549 *Environ Microbiol*. 56(11):3405-3411.
- 550 Yamane I, Sato K (1961) Effect of temperature on the formation of gases and ammonium nitrogen in the
- 551 water-logged soils. *Sci. Rep. Res. Inst. Tohoku Univ. D(Agr)*. 12:31-46.
- 552
- 553 The English in this document has been checked by at least two professional editors, both native speakers of
- 554 English. For a certificate, please see:
- 555
- 556 <http://www.textcheck.com/certificate/pm8wW1>
- 557
- 558

559 Figure1



560

561

562

563


564

565


566

567

568 Figure 2



581 Figure 3



582

583

584

585


586

587

588

589


590 Figure 4



591

592

593 Figure 5



594

595

596

597 Figure 6

